

Figure 1: Left panel: Crude numbers of LMS (dark grey bars) and non-LMS PCI (light grey bars) performed in England and Wales 2007-2014; Right panel: Percentage of total PCI performed in England and Wales 2007-2014 represented by LMS-PCI ($p < 0.001$ for trend).




Figure 2 Left panel: Patients undergoing LMS-PCI with an unprotected left main stem as a percentage of total LMS-PCI volume in England and Wales per year (p -value=0.696, $r^2=0.027$ for trend comparison); Right panel: Crude numbers of protected LMS-PCI (light grey bars) vs. unprotected LMS-PCI (dark grey bars) performed in England and Wales 2007-2014





Figure 3: Left panel - Temporal change in femoral access use for unprotected LMS-PCI in England and Wales 2007-2014 ($p < 0.001$ for trend); Right panel - Changes in left and right radial access for unprotected LMS-PCI in England and Wales 2007-2014 ($p < 0.001$ for both trends, $p < 0.001$ for comparison between trends).




Figure 4: Temporal changes in case complexity in the radial cohort of patients undergoing unprotected LMS-PCI in England and Wales 2007-2014 ($p < 0.001$ for all trends).




Figure 5: Temporal changes in event rates per year for the combined radial and femoral cohorts of patients undergoing unprotected LMS-PCI in England and Wales 2007-2014 ($p < 0.001$ for all trends).