

U–Pb dating of calcite in ancient carbonates for age estimates of syn- to post-depositional processes: a case study from the upper Ediacaran strata of Finnmark, Arctic Norway

Guido Meinhold^{1,2,*}, Nick M. W. Roberts³, Arzu Arslan¹, Sören Jensen⁴, Jan Ove R. Ebbestad⁵, Anette E. S. Högström⁶, Magne Høyberget⁷, Heda Agić⁸, Teodoro Palacios⁴ and Wendy L. Taylor⁹

¹School of Geography, Geology and the Environment, Keele University, Keele, ST5 5BG, UK;

²Department of Sedimentology and Environmental Geology, University of Göttingen, Goldschmidtstraße 3, D-37077 Göttingen, Germany;

³Geochronology and Tracers Facility, British Geological Survey, Nottingham, NG12 5GG, UK;

⁴Área de Paleontología, Facultad de Ciencias, Universidad de Extremadura, Avenida de Física, E-06006 Badajoz, Spain;

⁵Museum of Evolution, Uppsala University, Norbyvägen 16, SE 752 36 Uppsala, Sweden;

⁶Arctic University Museum of Norway, UiT - The Arctic University of Norway, N-9037 Tromsø, Norway;

⁷Rennesveien 14, N-4513 Mandal, Norway

⁸Department of Earth Science, University of California at Santa Barbara, Santa Barbara, CA 93106, USA and

⁹Department of Geological Sciences, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa

Author for correspondence:

Guido Meinhold, Email: g.meinhold@keele.ac.uk

Abstract

Results of *in situ* U–Pb dating of calcite spherulites, cone-in-cone (CIC) calcite and calcite fibres from a calcareous concretion of the upper Ediacaran of Finnmark, Arctic Norway, are reported. Calcite spherulites from the innermost layers of the concretion yielded a lower intercept age of 563 ± 70 Ma, which although imprecise, is within uncertainty of the age of sedimentation based on fossil assemblages. Non-deformed CIC calcite from the bottom part of the concretion yielded an age of 475 ± 25 Ma, which is interpreted as the age of CIC calcite formation during a period of fluid overpressure induced during burial of the sediments. Deformed CIC calcite from the top part of the concretion yielded an age of 418 ± 23 Ma, which overlaps with a known Caledonian tectono-metamorphic event, and indicates a potential post-depositional overprint at this time. Calcite fibres that grew in small fissures along spherulite rims, which are interpreted as a recrystallization feature during deformation and formation of a cleavage, gave an imprecise age of 486 ± 161 Ma. Our results show that U–Pb dating of calcite can provide age constraints for ancient carbonates and syn- to post-depositional processes that operated during burial and metamorphic overprinting.

Keywords:

U–Pb calcite geochronology; calcite spherulites; cone-in-cone calcite; Ediacaran; Caledonides; Norway

1. Introduction

Calcite U–Pb geochronology has attracted increasing interest in recent years within the Earth Sciences community. The method provides constraints on the age of sediment deposition and diagenesis (e.g. Israelson *et al.* 1996; Rasbury & Cole, 2009; Hill *et al.* 2016; Godeau *et al.* 2018; Pisapia *et al.* 2018; Drost *et al.* 2019), fossils (e.g. Rasbury & Cole, 2009; Yokoyama *et al.* 2018; Drost *et al.* 2019), and mineralization along fracture and fault planes, among others (e.g. Roberts & Walker, 2016; Goodfellow *et al.* 2017; Nuriel *et al.* 2017; Parrish *et al.* 2018; Holdsworth *et al.* 2019). Regardless of the successful application of calcite U–Pb geochronology in recent years, the method has its challenges. Calcite is typically low in U and rich in initial Pb, it is also susceptible to alteration or recrystallization at low temperature in the presence of fluids, and allows Pb diffusion above moderate temperatures (Cherniak, 1997). Carbonate formation can be complex and long-lived (Rasbury & Cole, 2009); the question therefore arises as to which geological ‘event’ is actually being dated (Rasbury & Cole, 2009; Drost *et al.* 2019; Roberts *et al.* 2020). The *in situ* technique, Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA–ICP–MS), helps overcome some of these challenges, namely by allowing for the measurement of discrete zones of uranium enrichment that are typical of diagenetic and hydrothermal calcite (Roberts *et al.* 2020), and by allowing a combination of U–Pb analysis with other *in situ* petrographic and analytical techniques.

In the present case study, we focus on calcite from carbonates of the upper Ediacaran–lower Cambrian Manndrapselva Member of the Stáhpogieddi Formation (Vestertana Group, Gaissa Nappe Complex) of the Digermulen Peninsula in eastern Finnmark, Arctic Norway (Fig. 1). The study area has attracted renewed research interest because of new findings of Ediacaran-aged fossils (e.g. Högström *et al.* 2013; Jensen *et al.* 2018a,b). The upper Ediacaran succession comprises siliciclastic sedimentary rocks with recently described carbonates, some with calcite spherulites and cone-in-cone (CIC) calcite (Meinhold *et al.* 2019a). Although the age of the carbonate-bearing part of the succession is well established as late Ediacaran based on fossils, the timing of the formation of the various types of calcite is poorly constrained. We therefore applied *in situ* U–Pb dating of calcite using LA–ICP–MS to address this question. Having different types of calcite in close proximity to each other, and having approximate age constraints of sedimentation based on biostratigraphy (Högström *et al.* 2013; McIlroy & Brasier, 2017; Jensen *et al.* 2018a,b) and of the low-grade metamorphic overprint (see discussion in Meinhold *et al.* 2019b), allows for the testing of the applicability of U–Pb calcite geochronology on a thin-section scale, and whether ages this far back into deep time can be related to a geological ‘event’ in a meaningful manner.

2. Geological setting

The study area is located in eastern Finnmark, Arctic Norway, and is part of the Gaissa Nappe Complex (Fig. 1a). The Stáhpogieddi Formation of the Vestertana Group has received much attention in recent years as it contains the only Ediacara-type fossils in Scandinavia as well as its most complete Ediacaran–Cambrian transition (Farmer *et al.* 1992; Högström *et al.* 2013; Jensen *et al.* 2018a,b) (Fig. 1b). The Manndrapselva Member of the Stáhpogieddi Formation consists of a basal sandstone-dominated part and two upwards-coarsening cycles. The trace fossil assemblage attests to its marine nature, and the sedimentology is consistent with deposition in a wave-dominated delta or shoreface (McIlroy & Brasier, 2017). The Ediacaran–Cambrian boundary is close to the base of the third cycle of the Manndrapselva Member based on trace fossils, palaeopascichnids and organic-walled microfossils (Högström *et al.* 2013; McIlroy & Brasier, 2017; Jensen *et al.* 2018a,b) (Fig. 1b).

Meinhold *et al.* (2019a) described carbonates within the second cycle of the Manndrapselva Member, which otherwise largely comprises alternating thin layers of silt- and mudstone and minor sandstone. Some of the sandstone beds show wave-formed ripple marks. Flute casts in the lower part of the succession indicate palaeocurrent flow from the N/NE. The rocks show cleavage, particularly pervasive in the muddy sediments.

The carbonates crop out along a coastal section at the eastern part of the Digermulen Peninsula (geographic coordinates: 70° 35' 31.0'' N, 28° 11' 30.3'' E) (Fig. 1c). They occur as beds, lenses, and concretions. Some consist of calcite spherulites and CIC structures made of calcite (see Meinhold *et al.* 2019a for details) (Fig. 2a, b). The upper Ediacaran sedimentary succession was deposited along the western margin of Baltica (in present-day coordinates) in a marine basinal environment (Fig. 1d). The rocks were metamorphosed during the Scandinavian Caledonian orogeny (Meinhold *et al.* 2019b).

3. Methodology

Bedrock sample material was cut with a rock saw perpendicular to the bedding to obtain a rock slice for thick-section preparation and *in situ* U–Pb dating of calcite (Fig. 2c–g). U–Pb geochronology of calcite was conducted at the Geochronology & Tracers Facility, British Geological Survey (Nottingham, UK), following the procedures described in Roberts & Walker (2016) and Roberts *et al.* (2017). The analyses were performed using a New Wave Research 193UC excimer laser ablation system, coupled to a Nu Instruments Attom single-collector sector-field ICP–MS. The method involves standard-sample bracketing with normalization to NIST 614 silicate glass (Woodhead & Hergt, 2001) for Pb–Pb ratios and WC1 carbonate reference material (Roberts *et al.* 2017) for U–Pb ratios. The laser parameters comprised a 80 µm static spot, fired at 10 Hz, with a ~3 J/cm² fluence, for 20 s of ablation. The material was pre-ablated to clean the sample site with 150 µm spots for 2 s. Data are plotted on a Tera-Wasserburg concordia diagram (²⁰⁷Pb/²⁰⁶Pb versus ²³⁸U/²⁰⁶Pb). The ages are determined by linear regression between common and radiogenic lead compositions and as lower intercepts on a Tera-Wasserburg concordia using the Microsoft Excel add-in Isoplot 4.15 (Ludwig, 2012). To keep track of precision and reproducibility of U–Pb ages, the Duff Brown Tank carbonate (²⁰⁶Pb/²³⁸U age = 64.04 ± 0.67 Ma; Hill *et al.* 2016) was analyzed in the course of this study. Measured isotopic ratios matched the published values of Hill *et al.* (2016) within uncertainty and yielded a lower intercept ²⁰⁶Pb/²³⁸U age of 66.3 ± 2.4 Ma (mean square weighted deviation (MSWD) = 7.8, N = 46). All ages are quoted at 2σ and include propagation of all systematic uncertainties (Horstwood *et al.* 2016). Full analytical data are provided in online Supplementary Table S1 and Figure S1 (available at <http://journals.cambridge.org/geo>).

4. Geochronological results

Four types of calcite from a carbonate concretion of the second cycle of the Manndrapselva Member of the Stáhpogieddi Formation were studied (Fig. 3). The domains include calcite spherulites, both undeformed and deformed CIC calcite, and calcite fibres grown in fissures along the spherulite rims (Fig. 2c–g). The majority of the ablated spots yielded low U and Pb contents ranging from less than 0.1 to 2.1 ppm (Average = 0.17 ppm, Median 0.06 ppm, N = 305) and 0.47 to 20.6 ppm (Average = 2.5 ppm, Median = 1.5 ppm, N = 305), respectively. The analyses of all samples were dominated by common lead, with only a small abundance of radiogenic lead, leading to large uncertainties on the regressed ages. The proportion of radiogenic lead varied in each sample, with the undeformed CIC calcite yielding the greatest abundance of radiogenic lead, and subsequently, the most precise age (Fig. 3b).

The following U–Pb lower intercept ages were obtained (Fig. 3). Calcite from calcite spherulites gave an age of 563 ± 70 Ma (MSWD = 1.11) (Fig. 3a). Calcite from the undeformed CIC calcite (bottom of concretion) gave an age of 475 ± 25 Ma (MSWD = 1.19) (Fig. 3b) whereas calcite from the deformed CIC calcite (top of concretion) yielded an age of 418 ± 23 Ma (MSWD = 0.67) (Fig. 3c). Calcite fibres grown in fissures along the spherulite rims gave an age of 486 ± 161 Ma (MSWD = 0.48) (Fig. 3d).

5. Discussion

The calcite spherulites are interpreted as a primary feature, forming in the sedimentary environment (Meinhold *et al.* 2019a). Within the large uncertainty (563 ± 70 Ma), the obtained age overlaps with the estimated timing of sedimentation (late Ediacaran; c. 545 Ma) based on body and trace fossil assemblages (Högström *et al.* 2013; McIlroy & Brasier, 2017; Jensen *et al.* 2018b) (Fig. 1b).

U–Pb ages from the CIC calcite from the bottom and the top of the concretion are surprisingly different. Undeformed CIC calcite (bottom of concretion) gave an age of 475 ± 25 Ma, whereas deformed CIC calcite (top of concretion) is younger, that is, 418 ± 23 Ma. The age of the undeformed CIC calcite is interpreted as the age of CIC calcite formation during a period of fluid overpressure as the sediments were buried. The age fits well with age estimates based on the required overburden to obtain the fluid overpressure needed to form CIC structures (see discussion in Meinhold *et al.* 2019a). In the case of the undeformed CIC calcite, the robust isochron (in terms of MSWD) implies that the isotopic system has remained a closed system (no loss or gain of U or Pb). The Early–Middle Ordovician age is interpreted as being meaningful and representing an approximate age estimate of the CIC formation. On the contrary, the apparent Silurian–Devonian age of the deformed CIC calcite is within the age range of the post-depositional overprint related to a late Caledonian tectono-metamorphic event in the Gaissa Nappe Complex of the Caledonides of Finnmark (see discussion in Meinhold *et al.* 2019b). This age is also robust in terms of MSWD (0.67) but lacks any measurement with abundant radiogenic lead. At face value, the regressed ages of the deformed and undeformed CIC are different, implying that the deformation of the CIC structures in the top part of the concretion may have occurred during the formation of the cleavage, and providing implications for the resetting of the U–Pb system in calcite.

Based on the colour of organic-walled microfossils from the Manndrapselva Member of the Stáhpogieddi Formation, the sedimentary rocks show a post-mature level indicating a thermal overprint of 200–250°C (T. Palacios, unpublished data, 2019). The maximum metamorphic overprint is given as low epizonal and reached around 300°C (see Meinhold *et al.* 2019b). If we assume that the CIC calcite from the top part of the concretion was originally undeformed and formed contemporaneously with the CIC calcite from the bottom part of the concretion (see Meinhold *et al.* 2019a), the data suggest that the original age of CIC calcite from the top part of the concretion has been reset during the Caledonian metamorphic overprint. Volume diffusion of Pb at temperatures of 250–300°C is a possibility, based on the experimental study of Pb diffusion in calcite (Cherniak, 1997), but mobility may also have been enhanced by grain deformation. Fluid infiltration is another possibility for resetting of the U–Pb system, although it is more likely to mobilize uranium since U(VI), in the form of uranyl ion (UO_2^{2+}), is highly soluble in oxidized waters (Langmuir, 1978); however, it may be expected that fluid-assisted alteration would have obliterated or at least affected the calcite growth structures, and this is not the case. We emphasize that although the data imply a resetting of the U–Pb system in the deformed CIC calcite, the lack of measured radiogenic lead, and hence precise age constraints, leads us to apply caution to this

interpretation. Nevertheless, the data provide compelling results that suggest the U–Pb calcite dating method has the potential for examining the timing of depositional, diagenetic and low-grade metamorphic events in sedimentary carbonates.

The calcite fibres grown in small fissures along the spherulite rims (Fig. 2c) gave an age of 486 ± 161 Ma, which is within uncertainty of both the sedimentation age, and the age of the metamorphic overprint and deformation. Formation of the calcite fibres is interpreted as being caused by dissolution and precipitation, during fracture and vein formation upon burial and compaction of the sediments. However, the uranium concentrations are extremely low, with the majority of analyses yielding < 0.0012 ppm, leading to a lack of measurable radiogenic lead and a large age uncertainty. We are therefore unable to refine the interpretation of the calcite fibres any further than that defined by the petrographic analysis (Meinhold *et al.* 2019a).

6. Conclusions

U–Pb calcite dates from an upper Ediacaran carbonate concretion provide timing constraints for depositional, diagenetic and potentially metamorphic processes, overlapping and confirming previous estimates based on relative bracketing of events. Our data show that LA–ICP–MS U–Pb dating of calcite can be a suitable complementary method to approximate the age of syn- to post-depositional processes that operate during burial and metamorphic overprinting and can be applied to ‘ancient’ carbonates. Of note, Precambrian sedimentary sequences often lack tight age constraints, particularly in settings where more robust geochronometers such as zircon (requiring cross-cutting intrusive or interbedded extrusive rocks) may be absent.

Acknowledgements. In the framework of the Digermulen Early Life Research Group, financial support for fieldwork in Arctic Norway was provided by the Research Council of Norway (grant No. 231103). We thank David Wilde for thick section preparation. Sören Jensen and Teodoro Palacios acknowledge funding from the Spanish ‘Ministerio de Economía, Industria y Competitividad’ (grant No. CGL 2017–87631P). We are very grateful to Bernard Bingen and Fernando Corfu for their supportive reviews.

Declaration of interest. None.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/xxxxxxx>

References

- Cherniak DJ** (1997) An experimental study of strontium and lead diffusion in calcite, and implications for carbonate diagenesis and metamorphism. *Geochimica et Cosmochimica Acta* **61**, 4173–79.
- Drost K, Chew D, Petrus JA, Scholze F, Woodhead JD, Schneider JW and Harper DAT** (2019) An image mapping approach to U–Pb LA–ICP–MS carbonate dating and applications to direct dating of carbonate sedimentation. *Geochemistry, Geophysics, Geosystems* **19**, 4631–48.
- Farmer J, Vidal G, Moczyłowska M, Strauss H, Ahlberg P and Siedlecka A** (1992) Ediacaran fossils from the Innerelv Member (late Proterozoic) of the Tanafjorden area, northeastern Finnmark. *Geological Magazine* **129**, 181–95.

- Godeau N, Deschamps P, Guihou A, Leonide P, Tendil A, Gerdes A, Hamelin B and Girard JP** (2018) U–Pb dating of calcite cement and diagenetic history in microporous carbonate reservoirs: Case of the Urgonian Limestone, France. *Geology* **46**, 247–50.
- Goodfellow BW, Viola G, Bingen B, Nuriel P and Kylander-Clark ARC** (2017) Palaeocene faulting in SE Sweden from U–Pb dating of slickenfibres calcite. *Terra Nova* **29**, 321–28.
- Hill CA, Polyak VJ, Asmerom Y and Provencio P** (2016) Constraints on a Late Cretaceous uplift, denudation, and incision of the Grand Canyon region, southwestern Colorado Plateau, USA, from U–Pb dating of lacustrine limestone. *Tectonics* **35**, 896–906.
- Holdsworth RE, McCaffrey KJW, Dempsey E, Roberts NMW, Hardman K, Morton A, Feely M, Hunt J, Conway A and Robertson A** (2019) Natural fracture propping and earthquake-induced oil migration in fractured basement reservoirs. *Geology*, doi: 10.1130/G46280.1.
- Horstwood MSA, Košler J, Gehrels G, Jackson SE, McLean NM, Paton C, Pearson NJ, Sircombe K, Sylvester P, Vermeesch P and Bowring JF** (2016) Community-derived standards for LA-ICP-MS U-(Th-)Pb geochronology – Uncertainty propagation, age interpretation and data reporting. *Geostandards and Geoanalytical Research* **40**, 311–32.
- Högström AES, Jensen S, Palacios T and Ebbestad JOR** (2013) New information on the Ediacaran–Cambrian transition in the Vestertana Group, Finnmark, northern Norway, from trace fossils and organic-walled microfossils. *Norwegian Journal of Geology* **93**, 95–106.
- Israelson C, Halliday AN and Buchardt B** (1996) U–Pb dating of calcite concretions from Cambrian black shales and the Phanerozoic time scale. *Earth and Planetary Science Letters* **141**, 153–9.
- Jensen S, Högström AES, Almond J, Taylor WL, Meinhold G, Høyberget M, Ebbestad JOR, Agić, H and Palacios T** (2018a) Scratch circles from the Ediacaran and Cambrian of Arctic Norway and the Republic of South Africa, with a review of scratch circle occurrences. *Bulletin of Geosciences* **93**, 287–304.
- Jensen S, Högström AES, Høyberget M, Meinhold G, McIlroy D, Ebbestad JOR, Taylor WL, Agić H and Palacios T** (2018b) New occurrences of *Palaeopascichnus* from the Stáhpogieddi Formation, Arctic Norway, and their bearing on the age of the Varanger Ice Age. *Canadian Journal of Earth Sciences* **55**, 1253–61.
- Langmuir D** (1978) Uranium solution-mineral equilibria at low temperatures with application to sedimentary ore deposits. *Geochimica et Cosmochimica Acta* **42**, 547–69.
- Ludwig K** (2012) User's Manual for Isoplot. A Geochronological Toolkit for Microsoft Excel. *Berkeley Geochronology Center, Special Publication no. 4*, 1–70.
- McIlroy D and Brasier MD** (2017) Ichneological evidence for the Cambrian explosion in the Ediacaran to Cambrian succession of Tanafjord, Finnmark, northern Norway. In *Earth System Evolution and Early Life: a Celebration of the Work of Martin Brasier* (eds AT Brasier, D McIlroy and N McLoughlin), pp. 351–68. Geological Society of London, Special Publication no. 488.
- Meert JG** (2014) Ediacaran–Ordovician paleomagnetism of Baltica: a review. *Gondwana Research* **25**, 159–69.
- Meinhold G, Jensen S, Høyberget M, Arslan A, Ebbestad JOR, Högström AES, Palacios T, Agić H and Taylor WL** (2019a) First record of carbonates with spherulites and cone-in-cone structures from the Precambrian of Arctic Norway, and their palaeoenvironmental significance. *Precambrian Research* **328**, 99–110.
- Meinhold G, Wemmer K, Högström AES, Ebbestad JOR, Jensen S, Palacios T, Høyberget M, Agić H and Taylor WL** (2019b) A late Caledonian tectono-thermal event in the Gaissa

- Nappe Complex, Arctic Norway: fine-fraction K–Ar evidence from the Digermulen Peninsula. *GFF* **141**, 289–94.
- Nuriel P, Weinberger R, Kylander-Clark ARC, Hacker BR and Craddock JP** (2017) The onset of the Dead Sea transform based on calcite age-strain analyses. *Geology* **45**, 587–90.
- Parrish RR, Parrish CM and Lasalle S** (2018) Vein calcite dating reveals Pyrenean orogen as cause of Paleogene deformation in southern England. *Journal of the Geological Society* **175**, 425–42.
- Pisapia C, Deschamps P, Battani A, Buschaert S, Guihou A, Hamelin B and Brulhet J** (2018) U/Pb dating of geodic calcite: new insights on Western Europe major tectonic events and associated diagenetic fluids. *Journal of the Geological Society* **175**, 60–70.
- Rasbury ET and Cole JM** (2009) Directly dating geologic events: U-Pb dating of carbonates. *Reviews of Geophysics* **47**, RG3001, doi: 10.1029/2007RG000246.
- Roberts NMW and Walker RJ** (2016) U-Pb geochronology of calcite mineralized faults; absolute dating of rift-related fault events on the northeast Atlantic margin. *Geology* **44**, 531–34.
- Roberts NMW, Rasbury ET, Parrish R, Smith CJ, Horstwood MSA and Condon DJ** (2017) A calcite reference material for LA-ICP-MS U-Pb geochronology. *Geochemistry, Geophysics, Geosystems* **18**, 2807–14.
- Roberts NMW, Drost K, Horstwood MSA, Condon DJ, Chew D, Drake H, Milodowski AE, McLean NM, Smye AJ, Walker RJ, Haslam R, Hodson K, Imber J, Beaudoin N and Lee JK** (2020) Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U–Pb carbonate geochronology: strategies, progress, and limitations. *Geochronology* **2**, 33–61.
- Siedlecka A, Reading HG, Williams GD and Roberts D** (2006) Langfjorden, preliminary bedrock geology map 2236 II, scale 1:50000. Norges Geologiske Undersøkelse, Trondheim.
- Woodhead JD and Hergt JM** (2001) Strontium, neodymium and lead isotope analyses of NIST glass certified reference materials: SRM 610, 612, 614. *Geostandards Newsletter* **25**, 261–6.
- Yokoyama T, Kimura J, Mitsuguchi T, Danhara T, Hirata T, Sakata S, Iwano H, Maruyama S, Chang Q, Miyazaki T, Murakami H and Saito-Kokubu Y** (2018) U-Pb dating of calcite using LA-ICP-MS: Instrumental setup for non-matrix-matched age dating and determination of analytical areas using elemental imaging. *Geochemical Journal* **52**, doi: 10.2343/geochemj.2.0541.




Fig. 1. (Colour online) (a) Outline of northernmost Scandinavia showing the Vestertana Group rocks, in grey shade, preserved within the Gaissa Nappe Complex (GNC) and parautochthonous in eastern Finnmark on the Varanger Peninsula (VP). Red box highlights the study area. TKFZ – Trollfjorden–Komagelva Fault Zone. (b) Simplified stratigraphy of the Vestertana Group (after Jensen *et al.* 2018b), showing occurrences of carbonates in the Ediacaran strata in eastern Finnmark. The stratigraphic position of carbonates, some with calcite spherulites and CIC calcite, analysed in the study are indicated 'CA'. (c) Geology of the SE portion of the Digermulen Peninsula, based on Siedlecka *et al.* (2006), showing locality where carbonates were found within the Mann-drapselva Member. We refer to Meinhold *et al.* (2019a) for details. (d) Late Ediacaran (550 Ma) palaeogeographic reconstruction of Baltica (after Meert, 2014). Land (ochre) and shallow sea (light blue) distributions were adopted from the palaeogeographic map series of Ron Blakey (Global Paleogeography and Tectonics in Deep Time ©2016 Colorado Plateau Geosystems Inc., used under an Academic Content License Agreement). Red box highlights the study area.




Fig. 2. (Colour online) (a) Field photograph showing calcite spherulites and undeformed CIC calcite from the second cycle of the Manndrapselva Member of the Stáhpogieddi Formation from the eastern part of the Digermulen Peninsula, Finnmark, Arctic Norway. (b) Schematic illustrations of calcite spherulite and CIC structures (after Meinhold *et al.* 2019a). (c–g) Images of the thick section from sample D17-GM4 used for *in situ* U–Pb dating of calcite. All images oriented with top up. (c) Entire thick section. Total length of the glass slide is 4.8 cm. The outer layer (top and bottom) consists of nested cones of fibrous calcite (CIC structures). The inner layers show thinly laminated calcareous siliciclastics and calcite spherulites. CIC structures of the bottom layer are undeformed whereas CIC structures of the top layer are deformed. Detailed descriptions are given in Meinhold *et al.* (2019a). Representative parts of the thick section studied by U–Pb geochronology are outlined with black frames. (d) Photomicrograph showing calcite spherulites with analysed spots (age group A). (e) Calcite spherulites with analysed spots (age group A) and calcite fibres grown in small fissures alongside the spherulite rims with analysed spots (age group D). (f) CIC structures of calcite from the outer layer (top) with analysed spots (age group C). (g) CIC structures of calcite from the outer layer (bottom) with analysed spots (age group B).




Fig. 3. (Colour online) *In situ* U–Pb dating of calcite. (a–d) Tera-Wasserburg concordia plots of $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{238}\text{U}/^{206}\text{Pb}$ ratios (uncorrected for common lead) for different types of calcite from sample D17-GM4 (second cycle of the Mandrapselva Member) measured *in situ* on a thick section by LA–ICP–MS. Sample details are given in Meinhold *et al.* (2019a). Each data-point ellipse denotes Pb/U ratios with error in 2σ uncertainty including propagation of systematic uncertainties for each laser-ablation spot. The lower intercept of the regression line through the majority of data indicates the age of calcite crystallization.